

University of Groningen

Perinatal tissue oxygenation and neurodevelopment in preterm and growth restricted infants

Richter, Anne

DOI:
[10.33612/diss.122713783](https://doi.org/10.33612/diss.122713783)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
Richter, A. (2020). *Perinatal tissue oxygenation and neurodevelopment in preterm and growth restricted infants*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen.
<https://doi.org/10.33612/diss.122713783>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Perinatal tissue oxygenation and neurodevelopment in preterm and growth restricted infants

Anne Elisabeth Richter

Perinatal tissue oxygenation and neurodevelopment in preterm and growth restricted infants.

Copyright © 2020 by A.E. Richter

Financial support for the printing of this thesis was kindly provided by the University of Groningen, University Medical Center Groningen, Graduate School of Medical Sciences, and the Research School of Behavioral and Cognitive Neurosciences.

ISBN: 978-94-034-2480-4

ISBN electronic version: 978-94-034-2479-8

Cover: A.E. Richter/Ridderprint BV, Ridderkerk, the Netherlands

Lay-out: A.E. Richter/Ridderprint BV, Ridderkerk, the Netherlands

Printed by: Ridderprint BV, Ridderkerk, the Netherlands



**rijksuniversiteit
 groningen**

Perinatal tissue oxygenation and neurodevelopment in preterm and growth restricted infants

Proefschrift

ter verkrijging van de graad van doctor aan de
Rijksuniversiteit Groningen
op gezag van de
rector magnificus prof. dr. C. Wijmenga
en volgens besluit van het College voor Promoties.

De openbare verdediging zal plaatsvinden op

maandag 20 april 2020 om 16.15 uur

door

Anne Elisabeth Richter

geboren op 14 december 1990
te Celle, Duitsland

Promotores

Prof. dr. A.F. Bos

Prof. dr. S.A. Scherjon

Copromotor

Dr. E.M.W. Kooi

Beoordelingscommissie

Prof. dr. A.A.E. Verhagen

Prof. dr. E. Lopriore

Prof. dr. E. Gratacos

Table of Contents

General introduction and outline of the thesis	7
Part I: Postnatal cerebral and systemic tissue oxygenation in relation to prenatal medication and placental dysfunction	35
Chapter 1: The effect of maternal antihypertensive drugs on the cerebral, renal, and splanchnic tissue oxygen extraction of preterm neonates <i>Neonatology 2016;110(3):163-171</i>	37
Chapter 2: Antenatal magnesium sulfate and preeclampsia differentially affect neonatal cerebral oxygenation <i>Provisionally accepted</i>	57
Chapter 3: Prenatal sildenafil and postnatal tissue oxygenation in fetal growth restriction <i>Submitted</i>	83
Part II: Outcomes associated with fetal brain-sparing and postnatal cerebral tissue oxygenation	107
Chapter 4: Postnatal cerebral hyperoxia is associated with an increased risk of severe retinopathy of prematurity <i>Neonatology 2019;116(4):356-362</i>	109
Chapter 5: Fetal brain-sparing, postnatal cerebral oxygenation, and neurodevelopment at 4 years of age following fetal growth restriction <i>Submitted</i>	127
Chapter 6: Altered neurodevelopmental DNA methylation status after fetal growth restriction with brain-sparing <i>Submitted</i>	159
General discussion and future perspectives	191
Appendix	227
English summary	229
Nederlandse samenvatting	235
Deutsche Zusammenfassung	243
Abbreviations	251
Dankwoord	253
About the author	259

General introduction and
outline of the thesis



8 | General introduction

‘Some 2.4 billion years ago, photosynthesis lead to the accumulation of oxygen to levels that were likely toxic to many obligate anaerobes. Organisms that could defend themselves against oxidative stress, and at the same time utilize oxygen for energy, survived and evolved. As time went on, a cellular requirement for oxygen became critical, and animals developed a biochemical response to low levels of oxygen.’

Sally L. Dunwoodie, The Role of Hypoxia in Development of the Mammalian Embryo¹

General introduction

The fetal circulation

Fetal cardiac development starts in the third week of gestation.² Innervation of the heart occurs at five weeks and by eight weeks it has developed into a four-chambered system resembling the adult heart.² Throughout gestation the heart continues to grow and adapt to increasing hemodynamic demands.³ Lung development starts at four weeks gestational age.⁴ The formation of sacculi from the terminal bronchioles with a first primitive air-blood-barrier, however, only begins at about 24 weeks gestational age, which limits effective ventilation of those born earlier.⁴ By 32 weeks, septation of the terminal sacs begins along with maturation of the surfactant system and capillary network and by 36 weeks the first true thin-walled alveoli responsible for effective gas exchange are visible.⁴

Until birth, the fetal lungs are filled with amniotic fluid and the fetus depends on oxygen and nutrient supply via the placenta. Therefore, the fetal circulation has to be different from the postnatal circulation. In short, highly oxygenated blood from the placenta is transported through the umbilical vein towards the liver. Half of the oxygenated blood is shunted away from the liver towards the heart through the ductus venosus.^{2,5} Although it mixes with de-oxygenated blood returning from fetal body parts into the vena cava, most of the oxygenated blood is directed towards the left atrium.^{2,5} From here, it follows the pressure gradient into the right atrium through a second shunt, the foramen ovale.^{2,5} Bypassing the right ventricle and lungs, well-oxygenated blood is thus directly shunted towards the left ventricle, which supplies the brain, heart, and upper limbs.^{2,5} Before descending further towards the lower fetal body parts, remaining blood mixes with less-oxygenated blood from the right ventricle, which has bypassed the pulmonary circulation through a third shunt, the ductus arteriosus, facilitated by a high pulmonary vascular resistance.^{2,5} A substantial fraction of the mixed blood

returns to the placenta via the umbilical arteries, while the remainder supplies the lower fetal body parts and abdominal viscera.^{5,6}

Fetal neurodevelopment

Brain development begins during week four of gestation with the formation of a forebrain (primary prosencephalon), midbrain (mesencephalon), and hindbrain (rhombencephalon) from the neural tube.⁷ The forebrain separates into the diencephalon and the secondary prosencephalon.⁸ While the former will give rise to the thalamus and epithalamus, the latter will develop into hypothalamus, retina, and telencephalon comprising cerebral cortex, hippocampus, and basal ganglia.^{8,9} The mesencephalon does not further subdivide but comprises the most cranial part of the brainstem.⁸ The medulla and pons of the brainstem and the cerebellum will develop from the hindbrain, from which also the spinal cord will extend.⁸

Brain development is initiated by rapid proliferation of neuronal precursor cells within the (periventricular) germinal matrix, which begin to differentiate at eight weeks gestation and subsequently migrate to their destined location.⁷ This results in growth and increased folding of the cortex.¹⁰ Following migration, which is completed at approximately 29 weeks gestation, the neurons start to synapse.⁷ A prominent role in this process plays brain-derived neurotrophic factor (BDNF), expression of which is dependent on neuronal activity.^{11,12} Established neural circuits are quickly stabilized by myelination of neuronal axons visible as white matter.⁷ Synaptogenesis and myelination continue long into adolescence and interruption can cause severe neurodevelopmental impairment.⁷

Vascularization of the brain

The brain derives its blood supply via the internal carotid and subclavian arteries, which gain their familiar morphology by seven weeks gestational age.² The first

branches into the anterior, middle, and posterior cerebral arteries (ACA, MCA, and PCA, respectively), which supply the cerebral hemispheres.¹³ The latter gives rise to the vertebral arteries, which join to become the basilar artery, supplying brainstem and cerebellum.¹³ A connection between the PCA and basilar artery develops, forming an anastomosis between the two vascular systems (the 'Circle of Willis'), while the carotid origin of the PCA regresses.^{14,15} Vascularization of the retina occurs only by the final trimester after maturation of the lens and regression of the hyaloid artery from lens to retina.^{9,16} This artery is later known as the central artery of the retina and constitutes a branch of the ophthalmic artery, which originates from the internal carotid.⁹

Intracerebral microvascularization develops early from the perineural (extracerebral) vascular plexus, from where capillaries penetrate deep into the brain tissue by angiogenic mechanisms, establishing an anastomotic plexus at the subventricular zone.¹⁷⁻¹⁹ From here, the brain parenchyma is vascularized in an ascending fashion, continuously reshaping according to neurodevelopmental needs, not to be complete within cortical areas until the age of 4 years.^{17,20,21} This angiogenic process is driven by vascular endothelial growth factor (VEGF), which also promotes survival of neurons and glial cells and axonal outgrowth.²² Interaction between the vascular endothelium and parenchymal astrocytes results in the formation of a blood-brain barrier (BBB), which limits the cerebral permeability to circulating cells or particles and safeguards cerebral extracellular homeostasis.²³ Maturation of the BBB is not complete until a few weeks after term.^{21,24}

An important feature of the brain is cerebrovascular autoregulation. It is the ability of the brain to preserve adequate cerebral perfusion despite fluctuations in blood pressure and oxygenation.²⁵ Functionality of adaptive vasodilation and constriction depends on the maturation of neurovascular mechanisms, which

starts at 23 weeks gestational age and is not yet fully established at term.^{19,21} Impairment of the cerebral autoregulation may cause cerebral ischemia and hemorrhage.²⁶

Fetal growth restriction

Fetal growth restriction (FGR) is a condition defined as fetal growth below its constitutional growth potential and is therefore not to be used interchangeably with being small-for-gestational age, which also includes fetuses or neonates being constitutionally small.²⁷ FGR may be caused by chromosomal or genetic syndromes, multiple gestation, congenital infection or metabolic disease, drug use, or inadequate maternal nutrition, but most common is placental insufficiency associated with gestational hypertension, preeclampsia (PE), diabetes, smoking, or other forms of abnormal uteroplacental development.^{28,29} Moreover, there seems to be a difference in placental pathology relative to onset of fetal growth restriction, with the placentas of early-onset fetal growth restriction (before approximately 32 weeks gestational age at diagnosis) showing more histologic signs of maternal vascular (uteroplacental) underperfusion, while late-onset fetal growth restriction is more associated with fetoplacental (villous) lesions.^{30,31}

Placental insufficiency and fetal hemodynamic redistribution

The placenta is the organ that supplies the fetus with oxygen and nutrients and removes waste back to the maternal circulation.³² Development of the placenta starts with the invasion of trophoblasts, the outer cell layer of the embryonic blastocyst, into the maternal endometrium and its capillary network.³³ This results in the formation of large sinuses, the so-called intervillous space, where exchange of gas and metabolites takes place.³³ The spiral arteries, which originate from the uterine arteries and supply the endometrium, are being remodeled and lose their smooth muscle layer, which turns them into highly dilated vessels with low

vascular resistance.³⁴ Fetal trophoblasts further bulge into the intervillous space to form the chorionic ('anchoring') villi, which develop an extensive network of capillaries connecting to the fetal umbilical veins.³⁵

Impaired remodeling of the spiral arteries is seen in early-onset PE.³⁶ Failure of dilation causes high jet velocities and high perfusion pressure in the intervillous space, which is damaging to chorionic villi and decreases the fetal capillary network.^{34,37} High uteroplacental resistance subsequently results in reduced placental perfusion and fetal hypoxia on the one hand and maternal hypertension and organ damage on the other.^{34,36} Although the pathophysiology of PE is much more complex and not yet fully understood, impaired placental perfusion can be detected during prenatal Doppler ultrasound as an increased resistance in the umbilical artery (UA), which normally decreases with progression of pregnancy.³⁸ As a compensatory response to hypoxia, the fetus redistributes its cardiac output with preferential perfusion of the brain at the expense of the systemic perfusion.³⁹ Whether this is neuroprotective and an active fetal mechanism is still controversial.^{40,41} However, this so-called 'brain-sparing' effect becomes apparent on prenatal Doppler as a decrease in resistance of the fetal MCA following cerebral vasodilation.⁴² Moreover, with progressive hypoxia, intracerebral regional redistribution seems to occur from higher towards lower order brain functions as the fetus starts to decompensate.⁴³⁻⁴⁵ Postnatally, body size of FGR infants with fetal brain-sparing may be asymmetric with a disproportionately large head (although still too small for gestational age) relative to body weight.⁴⁰

There has been much interest in the prevention and therapy of placental insufficiency. Promising options include oral phosphodiesterase 5 (PDE-5) inhibitors and local injections increasing placental VEGF expression.^{46,47} However, up to date, no effective and safe treatment of placental insufficiency exists and

FGR remains associated with a high risk of perinatal death and (iatrogenic) preterm birth.⁴⁷

Preterm birth

Preterm birth is defined as birth before 37 weeks gestational age and its complications are the leading cause of global childhood mortality under five years of age.⁴⁸ In Europe, about 9% of all live births are preterm and of these, 15% occur before 32 weeks' gestation.⁴⁹ Well known causes of preterm birth comprise placental insufficiency and fetal growth restriction, maternal respiratory and urinary infections, abnormal placentation, cervical insufficiency, ascending intrauterine infection, and/or congenital anomalies.⁵⁰⁻⁵² Further risk factors include multiple pregnancy, maternal stress and nutritional status, substance abuse, and genetic predisposition.⁵³ However, the majority of preterm births will be classified as idiopathic as no identifiable reason can be found.

Postnatal hemodynamics

With birth, the fetus enters a transitional phase, in which it adapts to postnatal life. It starts with the first breath and inflation of the lungs, which instantly lowers the pulmonary vascular resistance and increases pulmonary perfusion by the right ventricle.⁵⁴ Fetal lung fluid is absorbed and enters blood plasma increasing the circulating volume.⁵⁴ (Early) clamping of the umbilical cord results in a sudden decrease in preload and increase in afterload to which cardiac contractility and output have to adapt.^{55,56} Preload gradually increases with improved ventilation and increased left atrial return from the lungs.⁵⁷ Cardiac remodeling towards left ventricular predominance begins.⁵⁷ Within hours to days, the ductus arteriosus, foramen ovale, and ductus venosus close.^{58,59}

These sudden hemodynamic changes are particularly challenging for preterm and FGR neonates, which usually need ventilatory and sometimes hemodynamic

support. The neonatal lungs may struggle with the clearance of amniotic fluid and inflation due to immaturity of sodium channels, cesarean delivery, and surfactant deficiency.⁶⁰ The heart has a reduced ability to increase contractility and output to maintain adequate perfusion. This may be associated with an immature myocardium in the preterm infant or a decreased cardiac reserve following FGR-associated compensatory cardiac adaptation to intrauterine hypoxia.⁶¹⁻⁶⁴ Perinatal asphyxia (oxygen deprivation) as it may occur with placental insufficiency, may further depress cardiac function.⁶⁴ A depressed cortisol and catecholamine surge due to cesarean delivery and/or immature adrenal glands further contributes to postnatal hypotension and ineffective clearance of lung fluid.⁶⁰ Moreover, the ductus arteriosus may fail to close, causing decreased systemic perfusion and pulmonary hypertension as blood is now shunted into the opposite direction, namely from the aortic arch back into the pulmonary vasculature.⁵ While any fluctuations in perfusion pressure and oxygenation especially affect organs with less effective autoregulation, such as the intestines or kidneys, cerebral autoregulation should spare the brain.⁶⁵ However, cerebral autoregulation in preterm FGR neonates is frequently impaired, which relates to a combination of immaturity, asphyxia-associated vasoparalysis, and overriding brain-sparing mechanisms.^{66,67} Sickness and maternal or neonatal medication may further interfere with autoregulation and stable cerebral hemodynamics.⁶⁸⁻⁷¹

Outcomes associated with preterm birth and FGR

Neonatal complications of preterm birth and/or FGR are manifold and largely associated with organ immaturity and underlying pathology.⁷² They include respiratory distress, bronchopulmonary dysplasia (BPD), persistent pulmonary hypertension (PPHN), a patent ductus arteriosus (PDA), necrotizing enterocolitis (NEC), hypoglycemia, hyperbilirubinemia, anemia, sepsis, peri- and intraventricular hemorrhage (PIVH), periventricular leukomalacia (PVL), and

retinopathy of prematurity (ROP).⁷³⁻⁷⁶ In The Netherlands, the survival rate for neonates born before 24 weeks gestational age is less than 10%, but with active intervention from 24.0 weeks onwards rises to about 60% at 24.0 weeks and almost 100% by 32.0 weeks of gestation.⁷⁷ Although advanced perinatal care has led to improved outcomes of preterm infants, survival without major morbidity, in particular of those born extremely preterm, remains a challenge.⁷⁸⁻⁸⁰ Moreover, there is an increased risk for chronic kidney disease, metabolic syndrome, cardiovascular disease, and neurodevelopmental problems in later life, particularly in those born following FGR.⁸¹⁻⁸⁵

Brain injury and neurodevelopmental abnormalities after preterm birth and FGR

Preterm birth has been associated with altered regional brain volumes. Major risk factors are white matter injury, which presents a loss of myelinating oligodendrocytes associated with hypoxia- or inflammation-induced excitotoxicity, and FGR.^{86,87} Structural abnormalities associated with FGR likely reflect regional hemodynamic redistribution and include reduced cortical, hippocampal, and cerebellar volume, altered vascularization, delayed myelination, reduced synapsing in pre-frontal or limbic systems, and impaired neuronal migration.⁸⁸ Accordingly, preterm and FGR infants more often struggle with abnormal neurodevelopmental outcomes than term and appropriate-for-gestational age infants. These include cerebral palsy or minor motor impairments, epilepsy, sensory impairments, cognitive deficits, behavioral abnormalities, attention deficit hyperactivity disorder (ADHD), autism spectrum disorder (ASD), and language difficulties.⁸⁹⁻⁹¹ Impairments may become apparent as minor academic difficulties but can also be severely disabling.⁹²

Assessment of brain damage is therefore routinely performed in preterm and FGR infants. Serial cranial ultrasound can effectively detect PIVH and PVL (ischemia of the white matter at the watershed zones around the lateral

ventricles), while magnetic resonance imaging is better in detecting cerebellar hemorrhage or mild white matter injury.^{93,94} (Amplitude-integrated) electroencephalography can further aid in the prediction of white matter damage-associated neurological outcome and assessment of the variability and complexity of the early motor repertoire (general movements) may predict motor abnormalities and cognitive dysfunction.⁹⁵⁻⁹⁸

Hypoxic brain injury

As one may expect, hypoxia plays a pivotal role in brain injury. Hypoxemia disturbs mitochondrial adenosine triphosphate (ATP) generation and consequently ATP-dependent maintenance of an electrochemical gradient across the cell-membrane.⁹⁹ In addition to cellular edema, this causes decreased uptake of extracellular glutamate by astrocytes and increased glutamate release by neurons.⁹⁹ Excess glutamate in turn overactivates synaptic and extrasynaptic glutamate receptors, which causes excess calcium influx and induces pro-apoptotic pathways.¹⁰⁰ Moreover, excessive calcium within the mitochondria enhances mitochondrial generation of harmful reactive oxygen species (ROS).^{101,102} This production of ROS requires oxygen, which explains an additional oxidative tissue injury by ROS upon reperfusion.¹⁰¹ In addition, microglial overactivation by glutamate causes disruption of the BBB and neuroinflammation.¹⁰³ However, a glutamate imbalance seems to be particularly harmful to immature oligodendrocytes, which express higher levels of glutamate receptors than mature oligodendrocyte.^{104,105} Prenatal magnesium sulfate (MgSO₄) intravenously administered before imminent preterm birth has shown to protect the immature brain from excitotoxic injury by blockade of glutamate receptors, which has shown to effectively decrease the risk of cerebral palsy.^{100,106,107} Since moderate hypoxia is necessary for early embryonic and fetal (neuro)development, hypoxia may be especially harmful in advancing gestational

age where metabolic demands increase.¹⁰⁸ However, if oxygen saturations are too low in early gestation this may disturb critical developmental processes, such as neuronal migration and synaptogenesis as outlined earlier.¹⁰⁸

Hyperoxic brain injury

Hyperoxia has also demonstrated to be harmful to immature brain tissue and has been associated with reduced synaptic plasticity and myelination.¹⁰⁹ While the fetal brain develops well at relatively low arterial oxygen tensions of about 25 mm Hg, slowly increasing until term, oxygen tension quickly rises to about 70 mm Hg following birth.^{109,110} A consequential increase in ROS generation from an oxygen-rich environment cannot be effectively counterbalanced until maturation of the antioxidant defense in the third trimester.¹⁰⁹ Thereby hyperoxia is particularly harmful at low gestational age and preterm birth. As ROS react with various kinds of molecules, oxidative stress may damage DNA and mitochondrial membranes and eventually induces apoptosis.¹¹¹ Moreover, ROS production seems to be especially high in motor neurons and disturb adjacent astrocytic glutamate reuptake.¹¹² In addition, ROS cause lipid peroxidation, which disrupts the myelin sheath.^{109,113} Damage is potentiated, if hyperoxia is preceded by hypoxia-induced excitotoxic damage, which commonly occurs with placental insufficiency or immaturity-associated hemodynamic fluctuations.¹⁰⁹

Alterations in gene expression and methylation by hypo- and hyperoxia

In addition to direct cell injury, hypo- and hyperoxia may also interfere with the expression of genes important for normal maturation and neurodevelopment. In the brain, this has become particularly evident for retinal tissue, in which a sudden rise in oxygen tension following preterm birth results in downregulation of VEGF and angiogenesis.¹¹⁴ Reduced retinal vascularization in turn leads to retinal hypoxia as the retina matures, upon which uncontrolled upregulation of VEGF and

angiogenesis result in the formation of tortuous, leaky vessels, hemorrhage, and in severe cases detachment of the retina and blindness.¹¹⁴ In the postnatal period, arterial oxygen saturation (SaO₂) is therefore kept within a narrow moderately-high range to reduce hyperoxic exposure and the incidence of ROP without increasing the risk of hypoxia, NEC, and mortality.¹¹⁵

The transcription factor hypoxia-inducible factor 1 (HIF1) is a key regulator of the hypoxic response, although other factors such as cAMP response element binding protein (CREB) also appear to be involved.¹¹⁶ The alpha-subunit of HIF1 stabilizes in response to low oxygen concentrations, dimerizes with its beta-subunit and binds to the hypoxia-response element (HRE) in the promotor region of its target genes.¹¹⁶ The most popular target genes include *VEGF* and erythropoietin (*EPO*), which stimulate angiogenesis and the production of erythrocytes, respectively, alongside neurotrophic functions.^{22,117} In addition, hypoxia is able to induce epigenetic modifications, which act in concert with HIF1, and thereby long-term alterations in genetic expression, which has been implicated as an important component of embryogenesis, neurodevelopment, and disease.^{116,118,119} Indeed, several neurodevelopmental disorders, such as schizophrenia and autism, have been linked to hypoxia and the downregulation of VEGF and glutamate receptor units.¹²⁰ Moreover, experimental studies have shown that deletion of the HRE region in the VEGF gene causes motor neuron degeneration and that hyperoxia in neonatal mice reduces the expression of BDNF, underlining the regulatory role of oxygen in neurodevelopment.^{121,122}

Monitoring tissue oxygenation with near-infrared spectroscopy

Postnatal regional tissue oxygen saturation (rSO₂) can non-invasively be measured in the neonate using near-infrared spectroscopy (NIRS). Several devices exist, but here we will only address the INVOS device as used at our neonatal intensive care unit. Its technology utilizes tissue translucency to near-infrared light at a

wavelength spectrum of 730-810 nm, which is differentially absorbed by oxygenated and de-oxygenated hemoglobin and maximally separated from the absorption spectrum of water, preventing its interference.¹²³ Melanin, a pigment found in hair and skin, and bilirubin, a breakdown product of heme present in the skin with neonatal icterus, also well absorb light within the same wavelengths, which can lower rSO_2 measurements and produce outliers.¹²⁴ However, directional changes in rSO_2 remain visible, allowing for adequate trend monitoring.¹²⁵

Near-infrared light is delivered to the tissue by an emitting diode located within the sensor placed on the skin above the tissue of interest. Unabsorbed light is reflected and detected by two optodes located next to each other on the same sensor, as visualized in Figure 1. The signal returning to the optode situated closer to the light source is subtracted from the signal returning to the optode further away, thereby reducing signals from superficial tissue such as skin and bone.¹²⁶




Figure 1. Monitoring (cerebral) tissue oxygenation with near-infrared spectroscopy. Adopted from Zaleski et al.¹²⁷

In contrast to pulse oximetry, which subtracts non-pulsatile flow measurements to be able to measure only arterial oxygen saturation, NIRS measures reflections from the whole tissue microcirculation.¹²⁸ As the venous bed is largest, rSO_2 largely reflects venous oxygenation.¹²⁹ Simultaneous use of NIRS and pulse oximetry therefore allows calculation of the fractional tissue oxygen extraction (FTOE) using the formula $FTOE = (SaO_2 - rSO_2)/SaO_2$, which is an indicator of tissue ischemia, as it reflects the balance between oxygen delivery and consumption.¹³⁰

NIRS is becoming increasingly popular in neonatal intensive care, although its clinical use and interpretation remains subject of extensive research.¹³¹ Cerebral NIRS has demonstrated to be of particular value in preterm and FGR infants as it visualizes changes in cerebral perfusion associated with cardiorespiratory immaturity and transitional complications.¹³² Simultaneous monitoring of the splanchnic and renal tissue can be useful to early detect and interpret changes in global oxygenation status.¹³³ Figure 2 depicts the typical sites for NIRS monitoring in the neonate. Moreover, multisite NIRS can reflect fetal hemodynamic redistribution and brain-sparing associated with FGR, which may persist for several days after birth.¹³⁴

Cerebral oxygenation in the detection of brain injury

In newborn piglets, a cerebral rSO_2 below 35% has been associated with ischemic tissue damage and abnormal cerebral function, although this value may vary with different types of sensors.^{135,136} Likewise, a high cerebral FTOE, reflecting tissue hypoxia either due to increased cerebral perfusion or decreased cerebral metabolism, has been associated with a higher risk of PVL.¹³⁷ Its association with PIVH is less straightforward, as the pathophysiology of PIVH involves initial hyperperfusion and subsequent hypoperfusion.^{138,139} Assessing the correlation between cerebral rSO_2 and blood pressure can, however, detect impairment of

cerebral autoregulation, predict the risk of PIVH and PVL, and may aid in hemodynamic therapy.¹⁴⁰




Figure 2. Sensor locations in cerebral, splanchnic, and renal near-infrared spectroscopy (NIRS).

Although gestational age-associated reference values for cerebral rSO_2 and FTOE have been determined for the postnatal period and attempts have already been made to reduce the cerebral hypo- and hyperoxic exposure, optimal saturation ranges in relation to brain development and function remain unknown.¹⁴¹⁻¹⁴⁴ To effectively interpret and address cerebral oxygenation in preterm and FGR infants and improve neurologic outcome, we first need to better define and comprehend cerebral normoxia and how it relates to perinatal complications and therapies.

Aims and outline of this thesis

Given the high vulnerability of the brain and hemodynamic system in preterm and FGR neonates during the perinatal period, this thesis aims to contribute to an improved understanding of the perinatal cerebral and systemic tissue oxygenation, including fetal brain-sparing, and associated neurodevelopmental outcome in this patient population.

Part I of this thesis deals with the influence of prenatal medication and placental dysfunction on the cerebral and systemic tissue oxygenation of the preterm and/or FGR infant. The following research questions will be answered:

Chapter 1: Do maternal medications with antihypertensive properties, such as labetalol, nifedipine, and MgSO_4 affect the cerebral, renal, and splanchnic tissue oxygenation of preterm infants?

Chapter 2: Are changes in postnatal cerebral tissue oxygenation following prenatal MgSO_4 exposure related to changes in cerebral metabolism or perfusion and can they be differentiated from underlying hemodynamic effects of PE and fetal brain-sparing?

Chapter 3: Does prenatal sildenafil treatment for severe-early onset FGR affect postnatal cerebral and renal tissue oxygenation?

Part II of this thesis will address neurologic outcomes of fetal brain-sparing and postnatal cerebral tissue oxygen saturation. The following research questions will be answered:

Chapter 4: Is a prolonged duration of cerebral hyperoxia associated with the development of severe retinopathy of prematurity?

Chapter 5: How do fetal brain-sparing and postnatal cerebral oxygenation relate to intelligence, behavior, and executive functioning at 4 years of age in children born following FGR?

Chapter 6: Is fetal brain-sparing in FGR associated with differential oxygen-dependent methylation of neuroprotective genes at 4 years of age?

References

1. Dunwoodie SL. The role of hypoxia in development of the mammalian embryo. *Developmental cell*. 2009;17(6):755-773.
2. Moore K, Persaud T, Torchia M. Cardiovascular system. In: *The developing human: Clinically oriented embryology*. 11th ed. Elsevier; 2020:263-314.
3. Kiserud T, Acharya G. The fetal circulation. *Prenat Diagn*. 2004;24(13):1049-1059.
4. Smith LJ, McKay KO, van Asperen PP, Selvadurai H, Fitzgerald DA. Normal development of the lung and premature birth. *Paediatr Respir Rev*. 2010;11(3):135-142.
5. Finnemore A, Groves A. Physiology of the fetal and transitional circulation. *Semin Fetal Neonatal Med*. 2015;20(4):210-216.
6. Kiserud T. Physiology of the fetal circulation. *Semin Fetal Neonatal Med*. 2005;10(6):493-503.
7. Tau GZ, Peterson BS. Normal development of brain circuits. *Neuropsychopharmacology*. 2010;35(1):147-168.
8. Carlson B. Nervous system. In: *Human embryology and developmental biology*. 6th ed. Elsevier; 2019:205-241.
9. Carlson B. Sense organs. In: *Human embryology and developmental biology*. 6th ed. Elsevier; 2019:255-178.
10. del Toro D, Ruff T, Cederfjäll E, et al. Regulation of cerebral cortex folding by controlling neuronal migration via FLRT adhesion molecules. *Cell*. 2017;169(4):621-635. e16.
11. Gottmann K, Mittmann T, Lessmann V. BDNF signaling in the formation, maturation and plasticity of glutamatergic and GABAergic synapses. *Experimental brain research*. 2009;199(3-4):203-234.
12. Lu B, Nagappan G, Lu Y. BDNF and synaptic plasticity, cognitive function, and dysfunction. In: *Neurotrophic factors*. Springer; 2014:223-250.
13. Crossman A, Neary D. Vasculature of the central nervous system. In: *Neuroanatomy: An illustrated colour text*. 6th ed. Elsevier; 2020:60-67.
14. Bradac G. Aortic arch and origin of the brachiocephalic and cerebral arteries. In: *Applied cerebral angiography: Normal anatomy and vascular pathology*. 3rd ed. Springer; 2017:1-10.
15. Bradac G. Posterior cerebral artery. In: *Applied cerebral angiography: Normal anatomy and vascular pathology*. 2017th ed. Springer; 2017:115-123.
16. Schoenwolf G, Bleyl S, Brauer P, Francis-West P. Development of the eyes. In: *Larsen's human embryology*. 5th ed. Churchill Livingstone; 2015:488-500.
17. Ferrer I, Kaste M, Kalimo H. Vascular diseases. In: Love S, Louis D, Ellison D, eds. *Greenfield's neuropathology*. Vol 1. 8th ed. London: Hodder Arnold; 2008:121-240.
18. Marin-Padilla M. The human brain intracerebral microvascular system: Development and structure. *Front Neuroanat*. 2012;6:38. doi: 10.3389/fnana.2012.00038.
19. Frösen J, Joutel A. Smooth muscle cells of intracranial vessels: From development to

- disease. *Cardiovasc Res.* 2018;114(4):501-512.
20. Raybaud C. Normal and abnormal embryology and development of the intracranial vascular system. *Neurosurg Clin N Am.* 2010;21(3):399-426.
21. Pearce WJ. Fetal cerebrovascular maturation: Effects of hypoxia. *Semin Pediatr Neurol.* 2018;28:17-28.
22. Carmeliet P, Storkebaum E. Vascular and neuronal effects of VEGF in the nervous system: Implications for neurological disorders. *Semin Cell Dev Biol.* 2002;13(1):39-53.
23. Liebner S, Czupalla CJ, Wolburg H. Current concepts of blood-brain barrier development. *Int J Dev Biol.* 2011;55(4-5):467-476.
24. Marc T. Brain development and the immune system: An introduction to inflammatory and infectious diseases of the child's brain. In: *Handbook of clinical neurology.* Vol 112. Elsevier; 2013:1087-1089.
25. Rhee CJ, da Costa CS, Austin T, Brady KM, Czosnyka M, Lee JK. Neonatal cerebrovascular autoregulation. *Pediatr Res.* 2018;84(5):602-610.
26. Greisen G. Autoregulation of cerebral blood flow in newborn babies. *Early Hum Dev.* 2005;81(5):423-428.
27. Beune IM, Bloomfield FH, Ganzevoort W, et al. Consensus based definition of growth restriction in the newborn. *J Pediatr.* 2018;196:71-76. e1.
28. Galan HL. Protocol 40: Fetal growth restriction. In: Queenan J, Spong C, Lockwood CJ, eds. *Protocols for High-Risk pregnancies: An Evidence-Based approach.* 6th ed. Wiley-Blackwell; 2015:340-352.
29. Nardoza LMM, Caetano ACR, Zamarian ACP, et al. Fetal growth restriction: Current knowledge. *Arch Gynecol Obstet.* 2017;295(5):1061-1077.
30. Mifsud W, Sebire NJ. Placental pathology in early-onset and late-onset fetal growth restriction. *Fetal Diagn Ther.* 2014;36(2):117-128.
31. Savchev S, Figueras F, Sanz-Cortes M, et al. Evaluation of an optimal gestational age cut-off for the definition of early- and late-onset fetal growth restriction. *Fetal Diagn Ther.* 2014;36(2):99-105.
32. Burton GJ, Jauniaux E. What is the placenta? *Am J Obstet Gynecol.* 2015;213(4):S6-S8.
33. Nicholson R. An overview of development, function and diseases. In: *The placenta: Development, function and diseases.* Nova Science Publishers; 2013:1-30.
34. Burton G, Woods A, Jauniaux E, Kingdom J. Rheological and physiological consequences of conversion of the maternal spiral arteries for uteroplacental blood flow during human pregnancy. *Placenta.* 2009;30(6):473-482.
35. Henriksen T, Holme A, Horne H, Holm M, Michelsen T. Human placenta development, structure, and organization in relation to function and fetal development. In: Dutteroy A, Basak S, eds. *Human placental trophoblasts: Impact of maternal nutrition.* CRC Press; 2015:3-17.
36. Fisher SJ. Why is placentation abnormal in preeclampsia? *Am J Obstet Gynecol.* 2015;213(4):S115-S122.

37. Burton GJ, Jauniaux E. Pathophysiology of placental-derived fetal growth restriction. *Am J Obstet Gynecol.* 2018;218(2):S745-S761.
38. Gagnon R. Placental insufficiency and its consequences. *Eur J Obstet Gynecol Reprod Biol.* 2003;110:S99-S107.
39. Giussani DA. The fetal brain sparing response to hypoxia: Physiological mechanisms. *J Physiol.* 2016;594(5):1215-1230.
40. Giussani DA. The vulnerable developing brain. *Proc Natl Acad Sci U S A.* 2011;108(7):2641-2642.
41. Cohen E, Baerts W, van Bel F. Brain-sparing in intrauterine growth restriction: Considerations for the neonatologist. *Neonatology.* 2015;108(4):269-276.
42. Cruz-Martinez R, Figueras F. The role of doppler and placental screening. *Best Pract Res Clin Obstet Gynaecol.* 2009;23(6):845-855.
43. Figueroa-Diesel H, Hernandez-Andrade E, Acosta-Rojas R, Cabero L, Gratacos E. Doppler changes in the main fetal brain arteries at different stages of hemodynamic adaptation in severe intrauterine growth restriction. *Ultrasound Obstet Gynecol.* 2007;30(3):297-302.
44. Hernandez-Andrade E, Figueroa-Diesel H, Jansson T, Rangel-Nava H, Gratacos E. Changes in regional fetal cerebral blood flow perfusion in relation to hemodynamic deterioration in severely growth-restricted fetuses. *Ultrasound Obstet Gynecol.* 2008;32(1):71-76.
45. Rossi A, Romanello I, Forzano L, Cecchia M, Ambrosini G, Marches D. Assessment of fetal brain vascularization using three-dimensional power doppler ultrasound angiography in pregnancies affected by late-onset fetal growth restriction. *Global Journal of Medical Research.* 2014;14(1):Online ISSN: 2249-4618.
46. Paauw ND, Terstappen F, Ganzevoort W, Joles JA, Gremmels H, Lely AT. Sildenafil during pregnancy: A preclinical meta-analysis on fetal growth and maternal blood pressure. *Hypertension.* 2017;70(5):998-1006.
47. Groom KM, David AL. The role of aspirin, heparin, and other interventions in the prevention and treatment of fetal growth restriction. *Am J Obstet Gynecol.* 2018;218(2S):S829-S840.
48. Liu L, Oza S, Hogan D, et al. Global, regional, and national causes of under-5 mortality in 2000–15: An updated systematic analysis with implications for the sustainable development goals. *The Lancet.* 2016;388(10063):3027-3035.
49. Chawanpaiboon S, Vogel JP, Moller A, et al. Global, regional, and national estimates of levels of preterm birth in 2014: A systematic review and modelling analysis. *The Lancet Global Health.* 2019;7(1):e37-e46.
50. Dijkstra K, Janssen HC, Kuczynski E, Lockwood CJ. Cervical length in uncomplicated pregnancy: A study of sociodemographic predictors of cervical changes across gestation. *Am J Obstet Gynecol.* 1999;180(3):639-644.
51. Frey HA, Klebanoff MA. The epidemiology, etiology, and costs of preterm birth. *Semin Fetal Neonatal Med.* 2016;21(2):68-73.
52. Purisch SE, Gyamfi-Bannerman C. Epidemiology of preterm birth. *Semin Perinatol.* 2017;41(7):387-391.

53. Goldenberg R, Culhane J, Iams J, Romero R. Preterm birth 1: Epidemiology and causes of preterm birth. *Obstetric Anesthesia Digest*. 2009;29(1):6-7.
54. Hooper SB, Te Pas AB, Kitchen MJ. Respiratory transition in the newborn: A three-phase process. *Arch Dis Child Fetal Neonatal Ed*. 2016;101(3):F266-71.
55. Ersdal HL, Linde J, Mduma E, Auestad B, Perlman J. Neonatal outcome following cord clamping after onset of spontaneous respiration. *Pediatrics*. 2014;134(2):265-272.
56. Stenning FJ, Hooper SB, Kluckow M, et al. Transfusion or timing: The role of blood volume in delayed cord clamping during the cardiovascular transition at birth. *Frontiers in Pediatrics*. 2019;7. doi: 10.3389/fped.2019.00405.
57. Leone TA, Finer NN. Foetal adaptation at birth. *Current Paediatrics*. 2006;16(6):373-378.
58. Morton SU, Brodsky D. Fetal physiology and the transition to extrauterine life. *Clin Perinatol*. 2016;43(3):395-407.
59. Kiserud T. The ductus venosus. *Semin Perinatol*. 2001;25(1):11-20.
60. Hillman NH, Kallapur SG, Jobe AH. Physiology of transition from intrauterine to extrauterine life. *Clin Perinatol*. 2012;39(4):769-783.
61. Kluckow M, Evans N. Low systemic blood flow in the preterm infant. *Semin Neonatol*. 2001;6(1):75-84.
62. Leipälä JA, Boldt T, Turpeinen U, Vuolteenaho O, Fellman V. Cardiac hypertrophy and altered hemodynamic adaptation in growth-restricted preterm infants. *Pediatr Res*. 2003;53(6):989-993.
63. Crispi F, Hernandez-Andrade E, Pelsers MM, et al. Cardiac dysfunction and cell damage across clinical stages of severity in growth-restricted fetuses. *Am J Obstet Gynecol*. 2008;199(3). doi: 10.1016/j.ajog.2008.06.056.
64. Wu T, Azhibekov T, Seri I. Transitional hemodynamics in preterm neonates: Clinical relevance. *Pediatrics & Neonatology*. 2016;57(1):7-18.
65. Cerbo RM, Maragliano R, Pozzi M, et al. Global perfusion assessment and tissue oxygen saturation in preterm infants: Where are we? *Early Hum Dev*. 2013;89 Suppl 1:S44-S46.
66. Pryds O, Greisen G, Lou H, Friis-Hansen B. Vasoparalysis associated with brain damage in asphyxiated term infants. *J Pediatr*. 1990;117(1 Pt 1):119-125.
67. Cohen E, Baerts W, Caicedo Dorado A, Naulaers G, van Bel F, Lemmers PMA. Cerebrovascular autoregulation in preterm fetal growth restricted neonates. *Arch Dis Child Fetal Neonatal Ed*. 2018;104(5):F467-F472.
68. Soul JS, Hammer PE, Tsuji M, et al. Fluctuating pressure-passivity is common in the cerebral circulation of sick premature infants. *Pediatr Res*. 2007;61(4):467-473.
69. Schat TE, van der Laan ME, Schurink M, et al. Abdominal near-infrared spectroscopy in preterm infants: A comparison of splanchnic oxygen saturation measurements at two abdominal locations. *Early Hum Dev*. 2014;90(7):371-375.
70. Eriksen VR, Hahn GH, Greisen G. Dopamine therapy is associated with impaired cerebral autoregulation in preterm infants. *Acta Paediatrica*. 2014;103(12):1221-1226.

71. Caicedo A, Thewissen L, Naulaers G, Lemmers P, Van Bel F, Van Huffel S. Effect of maternal use of labetalol on the cerebral autoregulation in premature infants. In: *Oxygen transport to tissue XXXV*. Springer; 2013:105-111.
72. Basso O, Wilcox A. Mortality risk among preterm babies: Immaturity versus underlying pathology. *Epidemiology*. 2010;21(4):521-527.
73. Bird TM, Bronstein JM, Hall RW, Lowery CL, Nugent R, Mays GP. Late preterm infants: Birth outcomes and health care utilization in the first year. *Pediatrics*. 2010;126(2):e311-9.
74. Platt M. Outcomes in preterm infants. *Public Health*. 2014;128(5):399-403.
75. Abraham A, Rejiya C. Problems of prematurity. In: Bhattacharya N, Stubblefield P, eds. *Human fetal growth and development*. Springer; 2016:553-559.
76. Manuck TA, Rice MM, Bailit JL, et al. Preterm neonatal morbidity and mortality by gestational age: A contemporary cohort. *Am J Obstet Gynecol*. 2016;215(1).
77. Perined. Perinatale zorg in nederland 2016. <https://assets.perined.nl/docs/7935f9c6-eaac-4f59-a150-307ae04efa27.pdf>. Published 2018. Updated 2018. Accessed 09/23, 2019.
78. Groenendaal F, Termote J, Van Der Heide-Jalving M, Van Haastert I, De Vries L. Complications affecting preterm neonates from 1991 to 2006: What have we gained? *Acta Paediatrica*. 2010;99(3):354-358.
79. Stoll BJ, Hansen NI, Bell EF, et al. Trends in care practices, morbidity, and mortality of extremely preterm neonates, 1993-2012. *JAMA*. 2015;314(10):1039-1051.
80. Stensvold HJ, Klingenberg C, Stoen R, et al. Neonatal morbidity and 1-year survival of extremely preterm infants. *Pediatrics*. 2017;139(3).
81. Scott MN, Taylor HG, Fristad MA, et al. Behavior disorders in extremely preterm/extremely low birth weight children in kindergarten. *J Dev Behav Pediatr*. 2012;33(3):202-213.
82. Leviton A, Fichorova RN, O'Shea TM, et al. Two-hit model of brain damage in the very preterm newborn: Small for gestational age and postnatal systemic inflammation. *Pediatr Res*. 2013;73(3):362-370.
83. Lundequist A, Böhm B, Lagercrantz H, Forssberg H, Smedler A. Cognitive outcome varies in adolescents born preterm, depending on gestational age, intrauterine growth and neonatal complications. *Acta Paediatrica*. 2015;104(3):292-299.
84. Serenius F, Ewald U, Farooqi A, et al. Neurodevelopmental outcomes among extremely preterm infants 6.5 years after active perinatal care in sweden. *JAMA pediatrics*. 2016;170(10):954-963.
85. Stritzke A, Thomas S, Amin H, Fusch C, Lodha A. Renal consequences of preterm birth. *Mol Cell Pediatr*. 2017;4(1).
86. Thompson DK, Warfield SK, Carlin JB, et al. Perinatal risk factors altering regional brain structure in the preterm infant. *Brain*. 2006;130(3):667-677.
87. Alexander B, Kelly CE, Adamson C, et al. Changes in neonatal regional brain volume associated with preterm birth and perinatal factors. *Neuroimage*. 2019;185:654-663.
88. Miller SL, Huppi PS, Mallard C. The consequences of fetal growth restriction

- on brain structure and neurodevelopmental outcome. *J Physiol (Lond)*. 2016;594(4):807-823.
89. Allen MC. Neurodevelopmental outcomes of preterm infants. *Curr Opin Neurol*. 2008;21(2):123-128.
90. Buchmayer S, Johansson S, Johansson A, Hultman CM, Sparen P, Cnattingius S. Can association between preterm birth and autism be explained by maternal or neonatal morbidity? *Pediatrics*. 2009;124(5):e817-25.
91. Sucksdorff M, Lehtonen L, Chudal R, et al. Preterm birth and poor fetal growth as risk factors of attention-deficit/ hyperactivity disorder. *Pediatrics*. 2015;136(3):e599-608.
92. Twilhaar ES, de Kieviet JF, Aarnoudse-Moens CS, van Elburg RM, Oosterlaan J. Academic performance of children born preterm: A meta-analysis and meta-regression. *Arch Dis Child Fetal Neonatal Ed*. 2018;103(4):F322-F330.
93. Leijser LM, de Bruïne FT, Steggerda SJ, van der Grond J, Walther FJ, van Wezel-Meijler G. Brain imaging findings in very preterm infants throughout the neonatal period: Part I. incidences and evolution of lesions, comparison between ultrasound and MRI. *Early Hum Dev*. 2009;85(2):101-109.
94. Plaisier A, Raets MM, Ecury-Goossen GM, et al. Serial cranial ultrasonography or early MRI for detecting preterm brain injury? *Arch Dis Child Fetal Neonatal Ed*. 2015;100(4):F293-300.
95. Song J, Xu F, Wang L, et al. Early amplitude-integrated electroencephalography predicts brain injury and neurological outcome in very preterm infants. *Sci Rep*. 2015;5.
96. Bruggink JL, Van Braeckel KN, Bos AF. The early motor repertoire of children born preterm is associated with intelligence at school age. *Pediatrics*. 2010;125(6):e1356-63.
97. Einspieler C, Bos AF, Libertus ME, Marschik PB. The general movement assessment helps us to identify preterm infants at risk for cognitive dysfunction. *Front Psychol*. 2016;7. doi: 10.3389/fpsyg.2016.00406.
98. Peyton C, Yang E, Msall ME, et al. White matter injury and general movements in high-risk preterm infants. *AJNR Am J Neuroradiol*. 2017;38(1):162-169.
99. Pregolato S, Chakkarapani E, Isles AR, Luyt K. Glutamate transport and preterm brain injury. *Front Psychol*. 2019;10. doi: 10.3389/fpsyg.2019.00417.
100. Burd I, Welling J, Kannan G, Johnston MV. Excitotoxicity as a common mechanism for fetal neuronal injury with hypoxia and intrauterine inflammation. *Adv Pharmacol*. 2016;76:85-101.
101. Ten VS, Starkov A. Hypoxic-ischemic injury in the developing brain: The role of reactive oxygen species originating in mitochondria. *Neurol Res Int*. 2012;2012. doi: 10.1155/2012/542976.
102. Feno S, Butera G, Vecellio Reane D, Rizzuto R, Raffaello A. Crosstalk between calcium and ROS in pathophysiological conditions. *Oxid Med Cell Longev*. 2019;2019. doi: 10.1155/2019/9324018.
103. Ziemka-Nalecz M, Jaworska J, Zalewska T. Insights into the neuroinflammatory responses after neonatal hypoxia-ischemia. *Journal of Neuropathology & Experimental Neurology*. 2017;76(8):644-654.

104. Khwaja O, Volpe JJ. Pathogenesis of cerebral white matter injury of prematurity. *Arch Dis Child Fetal Neonatal Ed.* 2008;93(2):F153-61.
105. Matute C. Glutamate and ATP signalling in white matter pathology. *J Anat.* 2011;219(1):53-64.
106. Conde-Agudelo A, Romero R. Antenatal magnesium sulfate for the prevention of cerebral palsy in preterm infants less than 34 weeks' gestation: A systematic review and metaanalysis. *Am J Obstet Gynecol.* 2009;200(6):595-609.
107. Clerc P, Young CA, Bordt EA, Grigore AM, Fiskum G, Polster BM. Magnesium sulfate protects against the bioenergetic consequences of chronic glutamate receptor stimulation. *PLoS One.* 2013;8(11). doi: 10.1371/journal.pone.0079982.
108. Alves de Alencar Rocha AK, Allison BJ, Yawno T, et al. Early- versus late-onset fetal growth restriction differentially affects the development of the fetal sheep brain. *Dev Neurosci.* 2017;39(1-4):141-155.
109. Reich B, Hoeber D, Bendix I, Felderhoff-Mueser U. Hyperoxia and the immature brain. *Dev Neurosci.* 2016;38(5):311-330.
110. Vali P, Lakshminrusimha S. The fetus can teach us: Oxygen and the pulmonary vasculature. *Children (Basel).* 2017;4(8). doi: 10.3390/children4080067.
111. Torres-Cuevas I, Parra-Llorca A, Sánchez-Illana A, et al. Oxygen and oxidative stress in the perinatal period. *Redox biology.* 2017;12:674-681.
112. Rao SD, Yin HZ, Weiss JH. Disruption of glial glutamate transport by reactive oxygen species produced in motor neurons. *J Neurosci.* 2003;23(7):2627-2633.
113. Ravera S, Bartolucci M, Cuccarolo P, et al. Oxidative stress in myelin sheath: The other face of the extramitochondrial oxidative phosphorylation ability. *Free Radic Res.* 2015;49(9):1156-1164.
114. Smith LE. Pathogenesis of retinopathy of prematurity. *Growth Horm IGF Res.* 2004;14 Suppl A:S140-4.
115. Vento M. Oxygen supplementation in the neonatal period: Changing the paradigm. *Neonatology.* 2014;105(4):323-331.
116. Watson JA, Watson CJ, McCann A, Baugh J. Epigenetics: The epicenter of the hypoxic response. *Epigenetics.* 2010;5(4):293-296.
117. Buemi M, Cavallaro E, Floccari F, et al. Erythropoietin and the brain: From neurodevelopment to neuroprotection. *Clin Sci (Lond).* 2002;103(3):275-282.
118. Tsai Y, Wu K. Epigenetic regulation of hypoxia-responsive gene expression: Focusing on chromatin and DNA modifications. *International journal of cancer.* 2014;134(2):249-256.
119. Vaiserman AM, Koliada AK. Early-life adversity and long-term neurobehavioral outcomes: Epigenome as a bridge? *Hum Genomics.* 2017;11(1).
120. Howell KR, Armstrong J. Vascular endothelial growth factor (VEGF) in neurodevelopmental disorders. *Current Behavioral Neuroscience Reports.* 2017;4(4):299-308.
121. Oosthuyse B, Moons L, Storkebaum E, et al. Deletion of the hypoxia-response element in the vascular endothelial growth factor promoter causes motor neuron

- degeneration. *Nat Genet.* 2001;28(2):131-138.
122. Sengoku T, Murray K, Wilson M. Neonatal hyperoxia induces alterations in neurotrophin gene expression. *International Journal of Developmental Neuroscience.* 2016;48:31-37.
 123. Murkin JM, Arango M. Near-infrared spectroscopy as an index of brain and tissue oxygenation. *Br J Anaesth.* 2009;103 Suppl 1. doi: 10.1093/bja/aep299.
 124. Widman RA, Anderson A, Gonopolskiy O, inventors; Covidien LP SL, assignee. System and method of resolving outliers in NIRS cerebral oximetry. patent US8644901B2. 2014.
 125. Madsen PL, Skak C, Rasmussen A, Secher NH. Interference of cerebral near-infrared oximetry in patients with icterus. *Anesthesia & Analgesia.* 2000;90(2):489-493.
 126. Wolf M, Greisen G. Advances in near-infrared spectroscopy to study the brain of the preterm and term neonate. *Clin Perinatol.* 2009;36(4):807-834.
 127. Zaleski KL, Kussman BD. Near-infrared spectroscopy in pediatric congenital heart disease. *J Cardiothorac Vasc Anesth.* 2019;In press. doi: 10.1053/j.jvca.2019.08.048.
 128. Sood BG, McLaughlin K, Cortez J. Near-infrared spectroscopy: Applications in neonates. . 2015;20(3):164-172.
 129. Watzman HM, Kurth CD, Montenegro LM, Rome J, Steven JM, Nicolson SC. Arterial and venous contributions to near-infrared cerebral oximetry. *Anesthesiology: The Journal of the American Society of Anesthesiologists.* 2000;93(4):947-953.
 130. Korček P, Straňák Z, Širc J, Naulaers G. The role of near-infrared spectroscopy monitoring in preterm infants. *J Perinat.* 2017;37(10):1070-1077.
 131. da Costa CS, Greisen G, Austin T. Is near-infrared spectroscopy clinically useful in the preterm infant? *Arch Dis Child Fetal Neonatal Ed.* 2015;100(6):F558-61.
 132. Dix LML, van Bel F, Lemmers PMA. Monitoring cerebral oxygenation in neonates: An update. *Frontiers in pediatrics.* 2017;5. doi: 10.3389/fped.2017.00046.
 133. Mebius MJ, du Marchie Sarvaas, Gideon J, Wolthuis DW, et al. Near-infrared spectroscopy as a predictor of clinical deterioration: A case report of two infants with duct-dependent congenital heart disease. *BMC Pediatrics.* 2017;17(1). doi: 10.1186/s12887-017-0839-3.
 134. Tanis JC, Boelen MR, Schmitz DM, et al. Correlation between doppler flow patterns in growth-restricted fetuses and neonatal circulation. *Ultrasound Obstet Gynecol.* 2016;48(2):210-216.
 135. Kurth CD, McCann JC, Wu J, Miles L, Loepke AW. Cerebral oxygen saturation-time threshold for hypoxic-ischemic injury in piglets. *Anesth Analg.* 2009;108(4):1268-1277.
 136. Dix LM, van Bel F, Baerts W, Lemmers PM. Comparing near-infrared spectroscopy devices and their sensors for monitoring regional cerebral oxygen saturation in the neonate. *Pediatr Res.* 2013;74(5):557-563.
 137. Verhagen EA, Keating P, ter Horst HJ, Martijn A, Bos AF. Cerebral oxygen saturation and extraction in preterm infants with transient periventricular

- echodensities. *Pediatrics*. 2009;124(1):294-301.
138. Verhagen EA, ter Horst HJ, Keating P, Martijn A, Van Braeckel KN, Bos AF. Cerebral oxygenation in preterm infants with germinal matrix–intraventricular hemorrhages. *Stroke*. 2010;41(12):2901-2907.
 139. Alderliesten T, Lemmers PM, Smarius JJ, van de Vosse, René E, Baerts W, van Bel F. Cerebral oxygenation, extraction, and autoregulation in very preterm infants who develop peri-intraventricular hemorrhage. *J Pediatr*. 2013;162(4):698-704. e2.
 140. Vesoulis ZA, Mathur AM. Cerebral autoregulation, brain injury, and the transitioning premature infant. *Frontiers in pediatrics*. 2017;5. doi: 10.3389/fped.2017.00064.
 141. Pichler G, Binder C, Avian A, Beckenbach E, Schmölzer GM, Urlesberger B. Reference ranges for regional cerebral tissue oxygen saturation and fractional oxygen extraction in neonates during immediate transition after birth. *J Pediatr*. 2013;163(6):1558-1563.
 142. Alderliesten T, Dix L, Baerts W, et al. Reference values of regional cerebral oxygen saturation during the first 3 days of life in preterm neonates. *Pediatr Res*. 2016;79(1-1):55-64.
 143. Pellicer A, Greisen G, Benders M, et al. The SafeBoosC phase II randomised clinical trial: A treatment guideline for targeted near-infrared-derived cerebral tissue oxygenation versus standard treatment in extremely preterm infants. *Neonatology*. 2013;104(3):171-178.
 144. Riera J, Hyttel-Sorensen S, Bravo MC, et al. The SafeBoosC phase II clinical trial: An analysis of the interventions related with the oximeter readings. *Arch Dis Child Fetal Neonatal Ed*. 2016;101(4):F333-8.

PART I

POSTNATAL CEREBRAL AND SYSTEMIC TISSUE OXYGENATION IN RELATION TO PRENATAL MEDICATION AND PLACENTAL DYSFUNCTION

